

Transient Dynamical Processing in the Moth Macroglomerulus

Christopher L. Buckley and Thomas Nowotny

Centre for Computational Neuroscience and Robotics
University of Sussex
Brighton, UK

22nd June, 2009

Sensitivity and
Convergence

MGC Architecture
and Disinhibition

A transient
dynamical model

Future work

My Background

- ▶ Degree in Physics (Edinburgh University).
- ▶ Ph.D in Neural Network Theory (Southampton University).

Transient
Dynamical
Processing in the
Moth
Macroglomerulus

Sensitivity and
Convergence

MGC Architecture
and Disinhibition

A transient
dynamical model

Future work

My Remit

Focus on the properties of the response of a single glomerulus to a single pheromone component.

- ▶ Sensitivity of the MGC to low pheromone concentrations.
- ▶ Dynamic Range.
- ▶ Role of disinhibition.

Transient
Dynamical
Processing in the
Moth
Macroglomerulus

Sensitivity and
Convergence

MGC Architecture
and Disinhibition

A transient
dynamical model

Future work

This Talk

- ▶ Sensitivity and convergence.
- ▶ Disinhibition and the MGC architecture.
- ▶ A transient dynamical model.
- ▶ Future work.

Transient
Dynamical
Processing in the
Moth
Macroglomerulus

Sensitivity and
Convergence

MGC Architecture
and Disinhibition

A transient
dynamical model

Future work

Sensitivity and Convergence

Transient
Dynamical
Processing in the
Moth
Macroglomerulus



Sensitivity and
Convergence

MGC Architecture
and Disinhibition


A transient
dynamical model

Future work

- ▶ Behavioural response is sensitive to 300 molecular events on the antennae of *Bombyx mori* (Schneider, D. 1970).
- ▶ The cardiac rhythm is sensitive to 6 molecular events on the antennae of *Spodoptera littoralis* (Angioy, AM. 2003)?
- ▶ The convergence from ORN's onto the MGC is thought to explain this sensitivity (hyperacuity).
- ▶ Question: Can the convergence of the ORN's onto a single LN account for this sensitivity?

Sensitivity and Convergence

Transient
Dynamical
Processing in the
Moth
Macroglomerulus



Sensitivity and
Convergence

MGC Architecture
and Disinhibition


A transient
dynamical model

Future work

- ▶ Model the ORN's as a Bernoulli process.
- ▶ The probability of an ORN firing (state 1) in the absence of pheromone is $P_n = (1600/17000)$.
- ▶ The probability of an ORN firing in the presence of pheromone is $P_n + P_s = 1600/17000 + 300/17000$.
- ▶ The convergence rate from ORN's to LN's in the moth is about 1000.

Sensitivity and Convergence

Transient
Dynamical
Processing in the
Moth
Macroglomerulus



Sensitivity and
Convergence

MGC Architecture
and Disinhibition


A transient
dynamical model

Future work

- ▶ 50% behavioural response rate (Schneider, D. 1970).
- ▶ In 30 specimens not one responded in the absence of pheromone (Schneider, D. 1970).
- ▶ Consequently we want the number of false positives to be less than about 3%.

Sensitivity and Convergence

Transient
Dynamical
Processing in the
Moth
Macroglomerulus



Sensitivity and
Convergence

MGC Architecture
and Disinhibition


A transient
dynamical model

Future work

- ▶ This can almost explain a sensitivity to 300 events on the antennae.
- ▶ Greater sensitivity could be achieved by further integration at the level of the whole MGC.

MGC Architecture and Disinhibition

Transient
Dynamical
Processing in the
Moth
Macroglomerulus



Sensitivity and
Convergence

MGC Architecture
and Disinhibition

A transient
dynamical model

Future work

- ▶ PN's have a strong positive response to stimulation at the ORN's.
- ▶ However LN's only interact with inhibitory connections?

MGC Architecture and Disinhibition


Transient
Dynamical
Processing in the
Moth
Macroglomerulus

Sensitivity and
Convergence

MGC Architecture
and Disinhibition

A transient
dynamical model


Future work



- ▶ Consequently, some have modeled direct ORN-PN connections (Linster, C. et. al. 1996).
- ▶ However, these connections have never been experimentally observed.

MGC Architecture and Disinhibition

Transient
Dynamical
Processing in the
Moth
Macroglomerulus



Sensitivity and
Convergence

MGC Architecture
and Disinhibition

A transient
dynamical model

Future work

- ▶ Others have suggested an explicit disinhibition process through a feedforward layered system (Av-Ron, E. et. al. 1995).
- ▶ Consistent with the temporal delays between ORN and PN responses implying polysynaptic pathways.
- ▶ More layers could provide extra integration and perhaps more sensitivity to pheromone.

What kind of system is the MGC

Transient
Dynamical
Processing in the
Moth
Macroglomerulus

- ▶ We know little about its connectivity but explicit distinct layers have not been observed.
- ▶ There are most certainly recurrent connections (it's not feedforward).
- ▶ Question: Can a semi-random recurrent inhibitory network without explicit layers or ORN-PN connections produce both a strong PN response and sensitivity to pheromone?

Sensitivity and
Convergence

MGC Architecture
and Disinhibition

A transient
dynamical model


Future work

Other Main Assumption

- ▶ Only transient response to stimulation.

The Model

Transient
Dynamical
Processing in the
Moth
Macroglomerulus



Sensitivity and
Convergence

MGC Architecture
and Disinhibition


A transient
dynamical model

Future work

- ▶ Spiking network and a formally equivalent rate model.
- ▶ Assign connections probabilistically (no PN-LN connections).
- ▶ Use formal conditions on the rate model to guarantee only transient dynamics.

The Model

Transient
Dynamical
Processing in the
Moth
Macroglomerulus



Sensitivity and
Convergence

MGC Architecture
and Disinhibition


A transient
dynamical model

Future work

- ▶ This architecture has a strong resemblance to the liquid and echo state paradigms (Maas, W. et. al. 2002 and Jaeger, H. 2002).
- ▶ They have been suggested as good models of cortical microcolumns.

Some first results

Transient
Dynamical
Processing in the
Moth
Macroglomerulus



Sensitivity and
Convergence

MGC Architecture
and Disinhibition


A transient
dynamical model

Future work

- ▶ Rich LN responses because of transient dynamics.
- ▶ The PN response is excitatory with a delay and sometimes is preceded by a brief inhibitory period (*Manduca sexta*).

Mechanism

Transient
Dynamical
Processing in the
Moth
Macroglomerulus



Sensitivity and
Convergence

MGC Architecture
and Disinhibition


A transient
dynamical model

Future work

- ▶ The fraction of LN's that are connected to ORN's are all winners.
- ▶ Most LN's are inhibited by these winners.
- ▶ *Many* of losers outweigh the *few* winners disinhibiting the PN's.
- ▶ PN's that receive input from LN's that are synapsed by an ORN will have a brief initial inhibitory period.

Inferring Network Structure

Transient
Dynamical
Processing in the
Moth
Macroglomerulus



Sensitivity and
Convergence

MGC Architecture
and Disinhibition


A transient
dynamical model

Future work

- ▶ The rate description of the system allows an analytical purchase on the systems dynamics.
- ▶ For instance we can see that the periods of excitation and inhibition in the LN response is related to asymmetries in the MGC's connectivity.
- ▶ We maybe able to infer a lot about the MGC's connectivity by examining the detailed structure of the LN responses.

Critical Dynamics in the MGC

Transient
Dynamical
Processing in the
Moth
Macroglomerulus



Sensitivity and
Convergence

MGC Architecture
and Disinhibition

A transient
dynamical model

Future work

- ▶ Our system exhibits only transient dynamics.
- ▶ Too weak or too strong MGC connectivity results in unresponsive or complex cyclic (or chaotic) dynamics respectively.
- ▶ Systems poised between these two regimes are said to be critical.

Critical Dynamics in the MGC

Transient
Dynamical
Processing in the
Moth
Macroglomerulus

- ▶ Critical systems have been shown to have a number of desirable properties (Kinouchi et. al. 2006).
 - ▶ Sensitivity to input.
 - ▶ Increased dynamic range.
 - ▶ Rich computational properties.
- ▶ Kinouchi et al. have already suggested the olfactory bulb in vertebrates exhibits critical dynamics.
- ▶ We may be able to ascertain whether the MGC exhibits critical dynamics by looking at the distribution of sizes of the LN's responses to pheromone.

Sensitivity and
Convergence

MGC Architecture
and Disinhibition

A transient
dynamical model

Future work

Transient
Dynamical
Processing in the
Moth
Macroglomerulus

Sensitivity and
Convergence

MGC Architecture
and Disinhibition

A transient
dynamical model

Future work

Thank you!